

Molecular tectonics: Generation of grid and porous diamondoid coordination networks by calixarene based tectons

Ovsyannikov A., Ferlay S., Solovieva S., Antipin I., Konovalov A., Kyritsakas N., Hosseini M.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Combinations of neutral tetrakis meta-pyridyl appended calix[4]arene derivatives in 1,3-A conformation as tectons 3 (tetrakis-meta-pyridyl- tetrathiacalix[4]arene) and 4 (tetrakis-meta-pyridyltetrathiatetramercaptocalix[4]arene) with MX_2 , behaving as a neutral metallatecton, lead to the formation of 2- and 3-D neutral coordination networks. In all cases, the metal centre adopts a distorted octahedral geometry with the two anions occupying the two apical positions. For compound 3- CdX_2 ($X = Cl$ or Br), 4- $CdBr_2$ and 4- $FeCl_2$ similar porous diamondoid 3D networks, resulting from the tetrakis monodentate behaviour of the organic tectons, are formed. In marked contrast, for 4- CoX_2 ($X = Cl$ or Br), only two out of the four pyridyl units take part in the connectivity pattern leading thus to the formation of 2D grid type architectures. This journal is © the Partner Organisations 2014.

<http://dx.doi.org/10.1039/c3ce42305k>
